

	Bone spec. no.	bl (cm)	mass (g)	bl <sub>1gm</sub> (cm)	model	$L_{\text{birth}}$ (cm)	AL (cm)	ASM (years)	99%AL (cm)	AA (years)	AD (years)	$L_{\text{birth}}$ To AL	$L_{1\text{gm}}$ To $L_{\text{death}}$	MGR (g day <sup>-1</sup> )
<i>Dactylosaurus</i>	MB.R. 786	4.400	1239	0.970	average	0.913	4.479	1.476	4.434	9.149	9	0.220	0.208	0.672
	MB.R. 776.2	3.820	1075	1.216	average	1.210	5.444	3.236	5.390	13.548	5	0.222	0.318	0.882
<i>Anarosaurus</i>	Wijk 09-472	4.350	2175	1.704	average	1.706	4.409	0.351	4.365	6.331	6	0.387	0.392	1.030
	Wijk 07-70	4.400	2200	1.667	LGM	1.628	4.496	1.681	4.451	5.287	5	0.362	0.379	1.457
	Wijk 09-58	4.900	2450	1.491	average	1.389	5.002	1.470	4.951	5.310	5	0.278	0.304	2.306
aff. <i>N. pusillus</i>	SMNS 92125	1.810	1200	0.543	average	0.571	1.797	0.510	1.662	4.589	7	0.318	0.300	0.602
	SMNS 50372c	1.715	1130	0.461	average	0.500	1.838	0.123	1.820	13.418	6	0.272	0.269	0.183
<i>N. pusillus</i>	T 4178	1.750	1150	0.814	average	0.829	1.783	0.525	1.765	8.453	7	0.465	0.465	0.401
	T 4211	1.650	1085	1.039	average	0.360	1.658	0.676	1.643	10.789	8	0.217	0.630	0.278
<i>N. edwardsii</i>	T 4758	2.870	600	0.866	LGM	0.933	2.804	1.011	2.776	9.638	5	0.333	0.302	0.265
	phz 153	3.970	832	0.814	average	0.877	6.518	2.772	6.477	43.022	8	0.136	0.205	0.159
<i>Serpianosaurus</i>	T 4510	3.000	355	0.840	average	0.402	4.457	3.940	4.413	57.549	12–13	0.091	0.280	0.047
	T 119	2.130	252	0.377	average	0.152	2.185	3.223	2.163	15.824	10–11	0.064	0.177	0.044