

	Bone spec. no.	bl (cm)	mass (g)	bl _{1gm} (cm)	model	Asymp. mass	ASM / AA	ASM / AD	MGR / Asympt. mass (day ⁻¹)	MGR / mass (day ⁻¹)
<i>Dactylosaurus</i>	MB.R. 786	4.400	1239	0.970	average	2308.679	0.161	0.164	0.0003	0.0005
	MB.R. 776.2	3.820	1075	1.216	average	3118.347	0.239	0.647	0.0003	0.0008
<i>Anarosaurus</i>	Wijk 09-472	4.350	2175	1.704	average	2264.776	0.055	0.059	0.0005	0.0005
	Wijk 07-70	4.400	2200	1.667	LGM	2347.039	0.318	0.336	0.0006	0.0007
	Wijk 09-58	4.900	2450	1.491	average	2606.367	0.277	0.294	0.0009	0.0009
aff. <i>N. pusillus</i>	SMNS 92125	1.810	1200	0.543	average	920.651	0.111	0.073	0.0007	0.0005
	SMNS 50372c	1.715	1130	0.461	average	1376.688	0.009	0.021	0.0001	0.0002
<i>N. pusillus</i>	T 4178	1.750	1150	0.814	average	1215.601	0.062	0.075	0.0003	0.0003
	T 4211	1.650	1085	1.039	average	1087.671	0.063	0.085	0.0003	0.0003
<i>N. edwardsii</i>	T 4758	2.870	600	0.866	LGM	1215.601	0.105	0.202	0.0002	0.0004
	phz 153	3.970	832	0.814	average	1087.671	0.064	0.347	0.0002	0.0002
<i>Serpianosaurus</i>	T 4510	3.000	355	0.840	average	559.312	0.068	0.303–0.328	0.0001	0.0001
	T 119	2.130	252	0.377	average	355.000	0.204	0.293–0.322	0.0001	0.0002